








預防輪狀病毒

你真係識?



五寶龍



BB 1 至 3 個月大就會對黑白對比強烈的圖案有反應。¹

香港兒童常規**接種**計劃暫時未納入輪狀病毒疫苗²

► **保護** BB 6 週就可以開始預防輪狀病毒³

► **3 個月至 2 歲**為感染輪狀病毒的高峰期⁴

► 可引致 BB 發燒、嘔吐甚至脫水或休克^{4,5}

重要信息：

• 輪狀病毒疫苗不適用於對疫苗的任何成份過敏的嬰兒。對於免疫力受損、正在使用抑制免疫力療程、已感染 HIV、於 42 日內曾接受輸血或血產品的嬰兒，暫時沒有對上述人士使用該疫苗後的安全性及效用的數據。• 最常見之副作用包括腹瀉、嘔吐、發燒及中耳炎。請向醫生查詢有關細節，他們備有較詳盡的作用資料。• 該疫苗不能預防其他原因所引起的腹瀉。與所有疫苗一樣，該疫苗未必能為所有接種者提供保護。• 接受疫苗前，請先徵詢醫生之專業意見。

參考資料：

1. ZIMMERMANN, Anita, et al. Arquivos brasileiros de oftalmologia. 2019; 82(3): 173-175. 2. Family Health Service, Department of Health, Hong Kong. Schedule of Hong Kong Childhood Immunisation Program. Available at: https://www.fhs.gov.hk/english/main_ser/child_health/child_health_recommend.html [Access on 24 May 2021] 3. WHO. Rotavirus vaccine information sheet. Retrieved Jun 2, 2021, from https://www.who.int/vaccine_safety/initiative/tools/Rotavirus_vaccine_rates_information_sheet_0618.pdf 4. Muendo C, et al. BMC Pediatr. 2018;18(1):323. 5. US Centers for Disease Control and Prevention. Rotavirus. Symptoms. Available at: <https://www.cdc.gov/rotavirus/about/symptoms.html> [Access on 24 May 2021]



MSD

美國默沙東藥廠有限公司